

# The effects of mantle dynamics and atmospheric escape on the atmosphere of terrestrial planets.

Cédric Gillmann<sup>1</sup>, Paul Tackley<sup>1</sup>

<sup>1</sup>*Institute fuer Geophysik, Department of Earth Sciences, ETH Zurich, Switzerland  
cedric.gillmann@erdw.ethz.ch*

Mars, Venus and the Earth exhibit very diverse surface conditions as the result of the 4.5 billion years evolution that led to the planets we observe today. We propose to study the interaction between the atmosphere and the inner dynamics of terrestrial planets by modeling the long term evolution of Mars and Venus. This lets us investigate both the history of the surface conditions of these planetary bodies and the way mantle dynamics and atmospheric evolution are linked.

We generate histories of the state of the atmosphere of these planets as a function of time. We propose to use the recent advances due to observation and modeling to constrain possible evolutions of the atmosphere of terrestrial planets. The processes that are of interest here are atmospheric escape, both hydrodynamic (for early evolution) and non-thermal (during the late history of the planets) and mantle degassing. We model the effects of these processes.

In order to simulate the global evolution of the atmospheres of terrestrial planets, we need to estimate the fluxes of volatiles entering and leaving those atmospheres. The incoming flux is obtained by modeling the degassing of the planetary mantle. First, several studies modeling volcanic activity of terrestrial planets have been published (for Mars, [2], [6]). The second source for this data is the direct modeling of the mantle dynamics, adapting the advanced StagYY code developed by Tackley [5], [7]. Finally, in the case of Mars, we consider the observation of the surface of the planets and the study of different volcanic landforms as a way to constrain our numerical results [4].

The atmospheric escape is the main drain of volatiles considered in our model. The main phase evolution of the planet (post 4 Ga) is dominated by non thermal escape. We model the evolution of the outgoing volatile fluxes associated with this process based on present-day ASPERA (Analyzer of Space Plasma and Energetic Atoms) measurements [3] and the decrease of the Extreme UV (EUV) flux with time. Recent studies of the non-thermal escape fluxes provide a comparison [1]. Thermal escape is important early in the evolution of the planets when EUV flux was higher and the energy brought to the atmosphere by solar emissions was large enough to induce a global outflow of the atmosphere: hydrodynamic escape. By combining atmospheric escape, volcanic production and lava composition, we can first operate an integration backward in time, from the present day state of the atmosphere, to investigate its past. We are also able to obtain surface conditions and estimate surface temperatures.

The results we obtain allow us to propose different scenarios for the long term evolution of the atmospheres of Mars and Venus depending on parameters such as abundance of volatiles in the mantle and atmospheric escape efficiency. We show that the present-day atmosphere of Mars is likely to be constituted by a large part of volcanic gases. With a low  $CO_2$  concentration in the magma (150 ppm), present atmosphere is constructed of 50% of volcanic gases emitted since 3.7 billion years ago, which corresponds to an age of 1.9 to 2.3 Gyr. The variations of  $CO_2$  pressure over this period seem relatively low (50 mbars at most). This seems in line with the assumption of a heavy loss of volatiles during the first 500 Myr. Corresponding surface temperature variations are likely to be small (several Kelvin) and would not be responsible for periods of flowing liquid surface water by themselves. Water is, however, abundant on Mars during the whole 4 billion

years evolution (between 30% and 150% of the present day water) but is unlikely to reside in the atmosphere or in liquid form.

In the case of Venus atmospheric escape is much lower than on Mars during the late evolution, in particular when considering the high mass of the atmosphere. This is due to the larger gravity of the planet. Moreover, even early on  $CO_2$  escapes at a lower rate by hydrodynamic escape. Other species, such as water, however, are lost efficiently, which could be the cause of the dry state of the present day atmosphere. This is in agreement with calculated isotopic ratios for noble gases such as Neon and Argon.

## References

- [1] Barabash, S., A. Fedorov, R. Lundin, R., J.-A. Sauvaud, 2007: Martian atmospheric erosion rates, *Science*, 315, 501.
- [2] Breuer D., Spohn T. , 2006: Viscosity of the Martian mantle and its initial temperature: Constraints from crust formation history and the evolution of the magnetic field. *Planetary and Space Science*, 54, 153–169.
- [3] Fox, J.L. , 2004:  $CO_2$ + dissociative recombination: A source of thermal and nonthermal C on Mars, *J. Geophys. Res.*, 109, A08306.
- [4] Greeley, R., Schneid, B.D. , 1991: Magma Generation on Mars: Amounts, Rates, and Comparisons with Earth, Moon, and Venus. *Science*, Vol. 254. no. 5034, pp. 996 - 998.
- [5] Keller, T. and P. J. Tackley, 2009: Towards self-consistent modeling of the Martian dichotomy: The influence of low-degree convection on crustal thickness distribution, *Icarus* 202 (2), 429-443.
- [6] O'Neill, C., Lenardic, A., Jellinek, A.M. and Kiefer, W.S. , 2007: Melt propagation and volcanism in mantle convection simulations, with applications for Martian volcanic and atmospheric evolution. *Journal of Geophys. Res.*, vol. 112.
- [7] Tackley, P. J. , 2008: Modeling compressible mantle convection with large viscosity contrasts in a three-dimensional spherical shell using the yin-yang grid, *Phys. Earth Planet. Inter*, 171 (1-4), 7-18.




Figure 1: Evolution of the volcanic production on Mars for different models, from observation and numerical modeling. Primordial crust has been excluded from this figure.